




VACCINES. WORTH A SHOT.
Schedule or catch up on your adult or childhood immunizations today.
unitypoint.org/find-a-doctor



Follow all of ScienceDaily's latest research news and top science headlines!

Science News

from research organizations

Print Email Share

Flood risk for Iowa farmland

Date: April 6, 2022
Source: University of Iowa

Summary: Researchers have created a detailed set of maps examining the flood risk for all farmland in Iowa. The maps show 450,000 acres of crops in areas with a 50% chance of flooding, and losses from flooding averaging \$230 million annually.

Share: f t p in

RELATED TOPICS

Plants & Animals

- > Agriculture and Food
- > Food and Agriculture
- > Endangered Plants
- > Soil Types

Earth & Climate

- > Floods
- > Landslides
- > Natural Disasters
- > Water

As farmers ready for planting season, a new study examines the flood risk for all cropland in Iowa.

The study from IIHR-Hydrosience and Engineering at the University of Iowa is the first to detail the flood risk to farmland statewide. The researchers used flood maps developed at the Iowa Flood Center, and incorporated data from the Federal Emergency Management Agency (FEMA) and the U.S. Department of Agriculture to create the crop flood-risk analysis.

Among the main findings:

- Nearly 450,000 acres of Iowa farmland are located in a two-year flood return period, meaning there's a 50% chance the land will flood in a given year. That's less than 2% of the total farmable land analyzed in the study.
- Iowa agriculture sees crop losses, on average, of \$230 million a year due to farming that takes place in flood-prone areas.

The researchers also identified four watersheds as most vulnerable to flooding and crop losses: Middle Cedar in east-central Iowa, North Raccoon and South Skunk in central Iowa, and West Nishnabotna in southwest Iowa.

The new agricultural flood-risk maps developed by the IIHR researchers can be updated to reflect changes in climate; changes in land use, such as a shift in farming in a location; and changes to the landscape, such as the addition of a road or other infrastructure, to give a continuous picture of the flood potential for farmland across the state.

"It's a comprehensive approach to help create solutions with information that helps farmers take a clear-eyed look at their land and for policymakers and others to use as a starting point to determine how Iowa's landscape can be best used to reduce flooding," says Enes Yildirim, graduate research assistant at IIHR and the study's corresponding author.

The researchers analyzed nearly 25 million acres of agricultural land in Iowa and farming operations from 2016 to 2020 to classify the flood risk according to eight scenarios: 2-year, 5-year, 10-year, 25-year, 50-year, 100-year, 200-year, and 500-year return periods. Cropland located in a 2-year return period has a 50% chance of flooding in a given year; farmland in a 5-year return period has a 20% of flooding in a given year; while farmland in a 100-year return period has a 1% chance of flooding in a given year.

The researchers then incorporated flood maps from FEMA and the U.S. Army Corps of Engineers along with data from the USDA, including crop type, yields, costs and price, planting frequency, and a corn suitability rating, which indexes a farmland's productivity.

"We have taken all this information from federal agencies and have tailored it to create a more dynamic picture about the current agricultural flood risk in Iowa," says Ibrahim Demir, associate professor in civil and environmental engineering at Iowa and a study co-author.

Iowa has seen its fair share of flooding. Since 1953, 29 flood-related disaster declarations have been issued for the state, according to FEMA. Major, if not historic, flooding has occurred four times over the past decade and a half alone -- in 2008, 2014, 2016, and 2019.

The new maps seek to address objectively the flood stress points, by showing farmland that is prone to chronic flooding and has low productivity yields compared to other areas.

"We highlight the \$230 million in average annualized losses to show that there is farmland that is frequently exposed to floods and has a low corn suitability rating -- why not consider changing its use?" Yildirim says. "That, of course, would require further conversations, but you have to look at the costs and benefits of continuing to farm that land."

Policymakers also can entertain what to do with farmland that is prone to regular flooding but is highly productive. That is especially true for cropland in the West Nishnabotna region in southwest Iowa, the researchers found.

"The West Nishnabotna is a region that has a high corn suitability rating but also is exposed to regular flooding," Yildirim says. "So, it might need extra protection from flooding to maintain food production, such as building a levee, for example."

The researchers found rotating crops had a negligible impact on flood losses.

The study, "Agricultural flood vulnerability assessment and risk quantification in Iowa," was published online Feb. 26 in the journal *Science of the Total Environment*.

The University of Iowa and the Iowa Water Center funded the research.

Story Source:

Materials provided by **University of Iowa**. Original written by Richard C. Lewis. *Note: Content may be edited for style and length.*

Related Multimedia:

- Maps showing flood risk in Iowa

Journal Reference:

- Enes Yildirim, Ibrahim Demir. **Agricultural flood vulnerability assessment and risk quantification in Iowa.** *Science of The Total Environment*, 2022; 826: 154165 DOI: 10.1016/j.scitotenv.2022.154165

Cite This Page:

MLA APA Chicago

University of Iowa. "Flood risk for Iowa farmland." ScienceDaily. ScienceDaily, 6 April 2022.
<www.sciencedaily.com/releases/2022/04/220406101626.htm>.

Featured Content

from New Scientist

Earth's vital rivers captured in stunning drone and satellite images

April 6, 2022 — The world's rivers and their modern challenges are captured in dramatic photographs drawn from River, a new nature documentary that highlights the scale and intricacy of Earth's life-giving arteries.

We can still avert climate catastrophe -- but there is barely time

April 5, 2022 — The IPCC's latest report shows we have the tools to make fast cuts in emissions -- all that's missing is the political will.

Fabric conditioners reduce the release of microfibres in tumble dryers

April 6, 2022 — Tumble-drying clothes produces microfibre pollution, but this is almost halved by using a tumble dryer sheet and an anti-wrinkle fabric conditioner.

Visit *New Scientist* for more global science stories >>>

RELATED STORIES

Burrowing Critters Increase Risk of Levee Failure

Dec. 6, 2021 — When burrowing animals dig into earthen levees to make their homes, they can weaken the infrastructure to the point of collapse and localized flooding -- but most flood risk models don't take that ...

Sinking Land Will Exacerbate Flooding from Sea Level Rise in Bay Area

Mar. 7, 2018 — Hazard maps use estimated sea level rise due to climate change to determine flooding risk for today's shoreline, but don't take into account that some land is sinking. A precise study of subsidence ...

Flood Risk from American Rivers Is Greatly Underestimated

Feb. 28, 2018 — A groundbreaking new study has found that 41 million Americans are at risk from flooding rivers, which is more than three times the current estimate --- based on regulatory flood maps --- of 13 ...

Complex Causes of Maldives Flooding

Aug. 7, 2017 — The causes of coastal flooding in the Maldives are more complex than previously thought, according to a new study. Researchers examined wave and sea level data around historic flood events and found ...



Most Popular

this week

PLANTS & ANIMALS

Study Shows: Fish Can Calculate

Not All Dietary Fiber Is Created Equal: Cereal Fiber but Not Fruit or Vegetable Fibers Are Linked With Lower Inflammation

CRISPR and HIV: New Technique in Human Blood Unveils Potential Paths Toward Cure

EARTH & CLIMATE

Ozone May Be Heating the Planet More Than We Realize

New Polymer Membrane Tech Improves Efficiency of Carbon Dioxide Capture


'Freeze-Thaw Battery' Is Adept at Preserving Its Energy

FOSSILS & RUINS

Blue-Eyed Humans Have a Single, Common Ancestor

Boy or Girl? It's in the Father's Genes

Disbelief in Human Evolution Linked to Greater Prejudice and Racism



Strange & Offbeat

PLANTS & ANIMALS

High-Ranking Naked Mole-Rats Are More Resilient

Like College Roommates, Vampire Bats Bond When Randomly Paired

Squid Recorded Color-Matching Substrate for the First Time

EARTH & CLIMATE

Fruit Flies Adapt Activity to 'White Nights'

Flowers' Unseen Colors Can Help Ensure Pollination, Survival

Ancient Helium Leaking from Core Offers Clues to Earth's Formation

FOSSILS & RUINS

T. Rex's Short Arms May Have Lowered Risk of Bites During Feeding Frenzies

New Technology Solves Mystery of Respiration in Tetrahymena

Smells Like Ancient Society: Scientists Find Ways to Study and Reconstruct Past Societies

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

Email Newsletters
RSS Feeds

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

Facebook
Twitter
LinkedIn

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

Leave Feedback
Contact Us